

Some remarks on the Kronheimer-Mrowka classes of algebraic surfaces

Rogier Brussee*

alg-geom/9308003

1 Introduction

Recently Kronheimer and Mrowka have announced a very interesting result, which sheds new light on the Donaldson polynomials [K-M]. They find recurrence relations between the Donaldson polynomials, by finding relations between the polynomials and the minimal genus of a smooth real surface representing an homology class. To be more precise we need a definition.

For a simply connected 4-manifold X with odd $b_+ \geq 3$, we denote the $SU(2)$ polynomials on $H_0(X) \oplus H_2(X)$ by $q_k(X)$. X is called *simple* if we have

$$q_k(X)(pt^2, -) = 4q_{k-1}(X), \quad d = 4k - \frac{3}{2}(1 + b_+).$$

For simple 4-manifolds it is convenient to label the polynomials by their degree on $H_2(X)$ i.e. we define

$$q_d(X) = \begin{cases} q_k(X)|_{H_2(X)} & \text{if } d = 4k - \frac{3}{2}(1 + b_+), \\ q_k(X)(pt, -)|_{H_2(X)} & \text{if } d = 4k - 2 - \frac{3}{2}(1 + b_+), \\ 0 & \text{otherwise} \end{cases}$$

The *Donaldson series* is then the formal power series $q(X) = \sum_d q_d(X)/d!$.

Theorem 1.1. (Kronheimer, Mrowka) *For every simple 4-manifold X there exist a finite number of Kronheimer-Mrowka classes $K_1, \dots, K_p \in H^2(X)$ and non zero rational numbers $a_1, \dots, a_p$ such that*

$$(i) \quad q(X) = e^{Q/2} \sum_{i=1}^n a_i e^{K_i},$$

*Mathematisches Institut Universität Bayreuth, Universitäts Straße 30, D-98440 Bayreuth, Germany, e-mail:Brussee@btm8x1.mat.uni-bayreuth.de

- (ii) $K_i \equiv w_2(X) \pmod{2}$, for all $i = 1, \dots, p$,
- (iii) if $K_i \in \{K_1, \dots, K_p\}$ then $-K_i \in \{K_1, \dots, K_p\}$,
- (iv) for every homologically nontrivial connected real surface Σ with $\Sigma^2 \geq 0$ and every Kronheimer-Mrowka class K_i we have

$$2g(\Sigma) - 2 \geq \Sigma^2 + K_i \cdot \Sigma.$$

Here Q is the intersection form. The Kronheimer Mrowka classes will be abbreviated KM-classes and are called the basic classes in [K-M]. Condition (ii) is reminiscent of the Wu formula, whereas condition (iv) is similar to the adjunction formula for the genus of a smooth algebraic curve. This suggests that for complex surfaces, the KM-classes should be closely related to the canonical divisor K_X . Indeed in the examples and the conjectural expression for the Donaldson series of elliptic surfaces in the announcement, the KM-classes are of type $K_i = \alpha_i K_{\min} + \sum \beta_{ij} E_j$ where $K_{\min}$ is the canonical divisor of the minimal model, $E_1, \dots, E_l$ the (-1) -curves, α_i a rational number with $|\alpha_i| \leq 1$ and $\beta_{ij} = \pm 1$. A formulation which is a little less obvious but which generalises better, as we will see, is $K_i = C - D$ where C, D are divisors such that $K_X = C + D$ and such that a multiple is effective. Hence in these cases the canonical class is a KM-class which is extremal from an algebraic geometric point of view.

To formulate the general relationship, we recall that the effective cone of a complex surface $\text{NE}(X)$, is the positive rational cone in $\text{NS}(X)_{\mathbf{Q}} \subset H^2(X, \mathbf{Q})$ generated by effective divisors. Let $\overline{\text{NE}}(X)$ be its closure in the norm topology.

Theorem 1.2. *For every KM-class K_i on a simple simply connected algebraic surface X , there is a unique decomposition $K_X = C_i + D_i$ in $\mathbf{Z}$ -divisors $C_i, D_i \in \overline{\text{NE}}(X)$ such that $K_i = C_i - D_i$. In particular $K_i = K_X$ if and only if there is a smooth hyperplane section H such that $2g(H) - 2 = H^2 + K_i \cdot H$.*

The results in [Kro] seem to indicate that a KM-class and a hyperplane section H as in the theorem exist, at least when there is an $\omega \in H^0(K)$ such that for sufficiently large k , $q_k(\omega + \bar{\omega}) \neq 0$. For minimal surfaces of general type this would imply the invariance of the canonical class up to sign under orientation preserving self-diffeomorphisms.

Corollary 1.3. *Assume in addition that X is minimal and of general type, then $K_i^2 \leq K_X^2$ with equality if and only if $K_i = \pm K_X$.*

By the Lefschetz (1,1) theorem [G-H, p. 163], the algebraicity of the KM-classes is equivalent to the K_i being of type $(1,1)$, In fact this is what

we will prove, using that the Donaldson polynomials are of pure Hodge type as in [Br1]. Since we assume that $p_g > 0$ this shows that the lattice spanned by the K_i is a proper sublattice. Moreover since the K_i are defined by the differentiable structure, they are contained in the fixed lattice of the variation of Hodge structures defined by a family of complex structures on X . In favourable circumstances this should force the KM-classes to be in $H^2(X, \mathbf{Z}) \cap [K_{\min}, E_1, \dots, E_l]$.

Another implication of the theorem is that the KM-classes are trivial on $H^{0,2}(X) \subset H^2(X, \mathbf{C})$. Thus we get the following corollary.

Corollary 1.4. *If X is a simple and simply connected surface, then for all $\omega \in H^0(K_X)$ we have*

$$q(\omega + \bar{\omega}) = q_0 e^{\int \omega \wedge \bar{\omega}}$$

where q_0 is the Donaldson polynomial of degree 0

It would be rather interesting to understand this formula from an algebraic geometric point of view, possibly clarifying the role of the simpleness condition. Combining the corollary with O'Grady's non vanishing result, we see that $q_0 \neq 0$ if X is of general type, p_g is odd and $|K_{\min}|$, the linear system of the canonical class of the minimal model, contains a reduced curve [O'G, th. 2.4], [Li2, appendix], [Mor, th. 1].

In a similar vein, since the Neron-Severi group has a non degenerate intersection form, the K_i are determined by their intersection products with divisors. Thus we get

Corollary 1.5. *For a simple simply connected surface, the Donaldson series $q(X)$ is determined by $q(X)|_{\text{NS}(X)}$.*

The corollary says that by knowing the algebraic part of all Donaldson polynomials we can reconstruct the transcendental part as well, i.e. in the simple case, the polynomials defined by Jun-Li [Li1] contain as much information as the full polynomials. Moreover since $\overline{\text{NE}} \cap K_X - \overline{\text{NE}}$ is a bounded subset of $\text{NS}(X, \mathbf{Q})$, and the K_i are integral, at least in principle, we get an effective bound on the number of Kronheimer Mrowka classes, hence on the number of polynomials one has to compute in order to reconstruct the Donaldson series.

2 Proof of theorem 1.2 and corollary 1.3

We have to prove that the KM-classes are of type $(1, 1)$. Accepting this, the rest of the statement of theorem 1.2 and corollary 1.3 is a consequence of

property (ii), (iii), and (iv) of the KM-classes.

By property (ii) we can write $K_i = K_X - 2D$ for some $\mathbf{Z}$ -divisor D . For every very ample line bundle $\mathcal{O}(H)$, we choose a smooth connected hyperplane section H . Then we get

$$2g(H) - 2 = H^2 + H \cdot K_X \geq H^2 + H \cdot K_i = H^2 + H \cdot K_X - 2H \cdot D$$

i.e. $D \cdot H \geq 0$. Thus by the duality of the closure of the effective cone $\overline{\text{NE}}(X)$ and the nef cone, we conclude that $D \in \overline{\text{NE}}(X)$ [Wil, prop. 2.3]. We also have $C := (K_X + K_i)/2 \in \overline{\text{NE}}$ by property (iii). Rewriting we get $K_X = C + D$ and $K_i = C - D$ as claimed. Note that nothing is gained by applying the inequality to other smooth connected divisors C with $C^2 \geq 0$, since such divisors are nef. Finally note that $H \cdot K_i = 2g(H) - 2 - H^2 = H \cdot K_X$ if and only if $D = 0$, since $\overline{\text{NE}} \cap H^\perp = (0)$. This proves theorem 1.2 up to algebraicity.

Now assume temporarily that X is minimal and of general type. Write $K_i^2 = K_X^2 + 4(D^2 - K_X \cdot D)$. Since K_X is nef, we get $K_i^2 \leq K_X^2$ if $D^2 \leq 0$, with equality iff $D^2 = K \cdot D = 0$. The latter is equivalent to $D = 0$ by the Hodge index theorem. Interchanging K_i and $-K_i$ if necessary, we are left with the case $C^2 > 0$, $D^2 > 0$. Since $C, D \in \overline{\text{NE}}$, we have $C \cdot D > 0$ by the numerical connectedness of K_X [BPV, prop VII.6.1] (strictly speaking we need that C and D are effective, but the proof of [loc. cit] carries over without change). Thus $K_X^2 > D^2$. Then again by the Hodge index theorem we get $(K \cdot D)^2 \geq K^2 D^2 > (D^2)^2$, so $K_i^2 < K^2$. This proves the corollary.

To prove that the KM-classes are of type (1,1), we need a slight generalization of [Br1, prop. 3.1]. For simplicity we restrict ourselves to the $\text{SU}(2)$ case and a statement about Hodge types, but the proof can easily be modified to show that all $\text{SO}(3)$ polynomials $q_{L,k}(X)$ with $L \in \text{NS}(X)$ come form an algebraic cycle (cf. [loc. cit.]). Consider the Donaldson polynomial q_d as an element of $S^d H^2(X)$ endowed with its natural Hodge structure.

Lemma 2.1. *For every $d \geq 0$, the Donaldson polynomials q_d are pure of type (d, d) .*

Proof. We temporarily index the polynomials on $H_2(X)$ by $k = c_2$. By [Br1, prop. 3.1] the lemma is true for q_k with odd $k \gg 0$ and evaluated on 2 dimensional classes. Here sufficiently large means that the moduli space $\mathcal{M}_k(X)$ of stable bundles on X for generic polarisation H is generically smooth and reduced of the proper dimension, and that the lower moduli spaces $\mathcal{M}_{k'}(X)$ for $k' < k$ have sufficiently low dimension, so that we can

apply Morgan's comparison result [Mor]. For the general case we use stabilization.

By [M-O, th. 2.1.1], for every k we can find an l_0 , and $\epsilon_1, \dots, \epsilon_l$ sufficiently small such that for a generic polarisation on the $l \geq l_0$ -fold blow-up of type $H - \sum \epsilon_i E_i$, the moduli space $\mathcal{M}_{k+l}(\hat{X}(x_1, \dots, x_l))$ is generically smooth of the proper dimension and the lower moduli spaces $\mathcal{M}_{k'}(\hat{X})$ with $k' < k$, have sufficiently low dimension. Thus $q_{k+l}(\hat{X}(x_1, \dots, x_l))$ is pure of type $(d+4l, d+4l)$.

Now by the blow-up formula [F-M, th. 4.3.1], we have

$$q_k(X) = (-\tfrac{1}{2})^l q_{k+l}(\hat{X}(x_1, \dots, x_l))(E_1^4, \dots, E_l^4, -).$$

Clearly the exceptional divisors E_i are pure of type $(1, 1)$, hence $q_k(X)$ is pure of type (d, d) . Finally

$$q_k(X)(pt, -) = -\tfrac{1}{2} q_{k+1}(\hat{X}(x))(E^6, -),$$

by [F-M, below cor. 4.3.2] (without proof) or [Br2], and so we can use the same argument as above. □

The rest of the proof is now straightforward. Let

$$C(X) = e^{-Q/2} q(X) = \sum a_i e^{K_i}.$$

Then $C(X) = \sum C_d(X)$, where $C_d(X)$ is a homogeneous polynomial of pure type (d, d) , since both $\exp(-Q/2)$ and $q(X)$ is a sum of homogeneous polynomials of pure type. Write $K_i = \alpha_i + \beta_i + \bar{\alpha}_i$, with $\alpha_i \in H^{0,2}$, $\beta_i \in H^{1,1}$. Let $z \in H^{1,1}$ be variable and denote by $\frac{\partial}{\partial z}$ the directional derivative. Then we have

$$\frac{\partial^n C(X)}{\partial z^n} = \sum_{d=0}^{\infty} \frac{\partial^n C_d}{\partial z^n} = \sum a_i \langle \beta_i, z \rangle^n e^{K_i}.$$

Clearly $\frac{\partial^n C_d}{\partial z^n}$ has pure type $(d-n, d-n)$. Thus, upon restriction to $H^{0,2}(X)$ only the constant term contributes, and we get

$$\left. \frac{\partial^n C(X)}{\partial z^n} \right|_{H^{0,2}} = \text{constant} = \sum a_i \langle \beta_i, z \rangle^n e^{\bar{\alpha}_i}.$$

We conclude that

$$\sum_{\alpha_i \neq 0} a_i \langle \beta_i, z \rangle^n e^{\bar{\alpha}_i} = 0.$$

Since z and n are arbitrary we find that the sum must be empty, i.e. all KM-classes K_i are of type $(1, 1)$. □

References

- [BPV] Barth, W., Peters, C. and Van de Ven, A.: *Compact complex surfaces*. Heidelberg, Berlin, New York: Springer 1984.
- [Br1] Brussee, R.: *On the (-1) curve conjecture of Friedman and Morgan*, Inv. Math. to appear 1993, e-print alg-geom/9209001.
- [Br2] Brussee, R.: *Donaldson polynomials of connected sums*, in preparation.
- [F-M] Friedman, R. and Morgan, J.W.: *Smooth four manifolds and complex surfaces*. Ergebnisse der Mathematik und ihrer Grenzgebiete 3. Folge Heidelberg, Berlin, New York: Springer (to appear 1993).
- [G-H] Griffiths, P., and Harris, J.: *Principles of algebraic geometry*. New York Chichester Brisbane Toronto: John Wiley & Sons 1978.
- [Kro] Kroneimer, P.B.: *The genus minimizing property of algebraic curves*. Bull. A.M.S. **29** p. 63–69 July 1993.
- [K-M] Kronheimer P.B., and Mrowka T.S.: *Recurrence relations and asymptotics for four-manifold invariants*. research announcement submitted to Bull. Amer. Math. Soc. 1993.
- [Li1] Li, J.: *Algebraic geometric interpretation of Donaldson’s polynomial invariants*. J. Diff. Geom. **37** 417–466 (1993).
- [Li2] Li, J.: *Kodaira dimension of moduli spaces of vector bundles of surfaces*, preprint UCLA 1993.
- [Mor] Morgan, J.W.: *Comparison of the Donaldson polynomial invariants with their algebro-geometric analogues*. preprint Columbia university New York (1992).
- [M-O] Morgan, J., and O’Grady, K.: *The smooth classification of fake $K3$ ’s and similar surfaces*. preprint 1992.
- [O’G] O’Grady, K.: *Algebro-geometric analogues of Donaldson’s polynomials*. Inv. Math. **107** 351–359 (1992).
- [Wil] Wilson, P.: *Towards birational classification of algebraic varieties* Bull. London Math. Soc. **19** 1–48 (1987)